

Effect of Traditional and Truss Access Cavity Designs on Residual Bacterial Levels in Mandibular First Molars: An In Vitro Confocal Microscopy Study

Aim To quantitatively compare residual bacterial levels in the pulp chambers of mandibular first molars prepared using traditional endodontic access cavities (TEC) and truss access cavities (TAC) following standardized cleaning and shaping procedures.

Methodology This in vitro experimental study included thirty extracted human mandibular first molars, randomly assigned to two groups ($n = 15$ each): TEC and TAC. After access cavity preparation, all specimens were inoculated with *Actinomyces naeslundii* and incubated for 10 days to allow biofilm formation. The distal canals were instrumented using Vortex Blue rotary files with continuous irrigation using 5.25% sodium hypochlorite at a flow rate of 1 mL/min. A final irrigation protocol consisting of 5 mL of 5.25% sodium hypochlorite, 17% ethylenediaminetetraacetic acid, and 5 mL of sterile saline was applied. Specimens were sectioned and analyzed using confocal laser scanning microscopy to quantify residual bacterial presence. Percentage bacterial reduction was calculated, and intergroup comparisons were performed using an independent two-sample t test ($\alpha = 0.05$).

Results The mean (\pm SD) percentage of bacterial reduction was $32.7\% \pm 8.0$ in the TEC group and $28.4\% \pm 4.8$ in the TAC group. No statistically significant difference was observed between the groups ($P = 0.08$).

Conclusion Under the conditions of this in vitro study, truss access cavities demonstrated bacterial reduction comparable to traditional endodontic access cavities following cleaning and shaping. These findings suggest that truss access designs may be considered a minimally invasive alternative without compromising intracanal disinfection.

Kapil Jadhav^{1*}
Cinnamon VanPutte²
Shiva Kumar³
Ajay Praveen⁴
Suresh Shenvi⁵
Kavalipurapu Venkata Teja⁶
Omer Sheriff Sultan⁷
Kaligotla Apoorva Vasundhara⁸
Vittoria Rodriquez⁹
Salvatore Scolavino⁹

¹Assistant Professor, A.T. Still University - Missouri School of Dentistry and Oral Health, St. Louis, MO

²Associate Professor, Southern Illinois University School of Dental Medicine, Alton, IL

³The Department of Forensic Odontology, Government Dental College, Research Institute, Bangalore, India

⁴Department of Conservative Dentistry and Endodontics, Government Dental College, Pudukkottai, India

⁵Department of Conservative Dentistry & Endodontics, KLE VK Institute of Dental Sciences, KLE Academy of Higher Education and Research, Deemed-to-be-University, Belagavi, Karnataka, India.

⁶Department of Conservative Dentistry and Endodontics, Malla Reddy Institute of Dental Sciences, Malla Reddy Vishwavidyapeeth, Hyderabad, Telangana, India.

⁷Associate Professor, A.T. Still University - Missouri School of Dentistry and Oral Health, St. Louis, MO

⁸Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India.

⁹Department of Neuroscience, Reproductive Sciences and Dentistry, University of Naples Federico II, Naples, Italy.

*Corresponding author

Received 2025, December 19

Accepted 2026, March 20

KEYWORDS Access designs, Dental pulp, Endodontics, Molars, Root canal.

Corresponding Authors*

Kapil Jadhav, Assistant Professor, A.T. Still University - Missouri School of Dentistry and Oral Health, St. Louis, MO | kapiljadhav@atsu.edu |

Peer review under responsibility of Società Italiana di Endodonzia

10.32067/GIE.2026.546

Società Italiana di Endodonzia. Production and hosting by Helyx srl on behalf of Tecniche Nuove. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The primary objective of root canal treatment is to eliminate harmful pathogens from the complex root canal system to preserve teeth and restore their function in the oral cavity (1). However, a combination of structural and microbiological factors may lead to failure of endodontically treated teeth (ETT) (2, 3). Structural factors can include loss of tooth structure from extensive caries or size of access cavity preparation, excess removal of radicular dentin during instrumentation, improper selection of post, or proteolytic and demineralizing actions of root canal irrigants (4-6). Microbiological factors can include incomplete debridement of pulpal tissues and remaining dentinal debris or viable microorganisms (7).

Historically, successful endodontic treatment has emphasized the complete removal of caries and the importance of root canal disinfection. Traditional endodontic access cavities (TEC) are commonly used in endodontic treatment and often result in premature, structural failure-derived loss of ETT, but highly flexible heat-treated rotary instruments and enhanced illumination and imaging technology have decreased the overall failure rate (8,9). Recently, the concept of minimally invasive endodontics has been suggested to increase the success of ETT, promote the preservation of hard dental tissues during root canal treatment, and improve longevity and functionality (10). Despite ongoing debate about the size of the access cavity and root canal preparation, no consensus exists regarding the best approach to ETT (11). Endodontic therapy involves cleaning and shaping, disinfection, and 3-dimensional obturation of the root canal system. During root canal treatment, the goal of an access cavity design is to obtain straight-line access for complete debridement of pulp and to enhance the cleaning and shaping of the root

canal system (8). When TEC design is used, the roof of the pulp chamber is completely removed to improve access to the coronal part of the root canal system (12). This process frequently results in removal of tooth structure, which weakens the tooth in the pericervical area, and predisposes it to fractures, which may cause the tooth to be nonrestorable (9-11).

Minimally invasive access cavity preparations, also known as conservative endodontic access cavity designs (CEC), are intended to preserve the pericervical dentin by retaining as much of the pulp chamber roof as possible with the aim of improving the fracture resistance of ETT (13). Truss access cavity (TAC) is a type of CEC that involves creating separate access cavities to treat each canal individually. For example, with mandibular molars, one access cavity is made for the mesial canals and another for the distal canals (14). The primary goal of TEC is to maximize dentin preservation by leaving a truss of dentin between access cavities, thus improving the fracture resistance of ETT (13-15). Improper post selection is a well-documented reason for the failure of ETT (16,17), so a benefit of using TAC is a decreased need for post placement for restoring ETT. In addition, TAC has been reported as less time consuming than TEC (18). However, if the access cavity is too small, it can be challenging to locate the orifices of canals. As such, it can be difficult to carry out the necessary cleaning, shaping, and obturation of the root canal system and may lead to endodontic mishaps like transportation and deviation of root canals (6,10,11,18-21). The importance of preserving dentin cannot be overemphasized and must be balanced with disinfecting abilities to achieve successful endodontic treatment. To date, studies have focused on the fracture resistance of ETT with various types of CEC, but there is limited data about the debridement of the root canals using CEC (14,21-24).

Endodontic infections are typically



polymicrobial in nature and often involve several microbial species that work synergistically to cause infection and progression of pulpal disease (25). Actinomycetes bacteria are anaerobic, Gram-positive rods that are frequently seen in primary and secondary endodontic infections. Actinomycetes work synergistically with Pseudopropionibacterium species, leading to persistent extraradicular infections (25,26). Both bacteria are members of the order of Actinomycetota and have the ability to colonize on external root surfaces. This colonization is the primary cause of persistent endodontic lesions, even in the absence of endodontic infection. In particular, the presence of *Actinomyces naeslundii* in a root canal system has been correlated with failure of endodontic treatment since this bacterium is difficult to eradicate by chemomechanical debridement (27,28). However, a study by Radcliffe et al. (25) reported several concentrations of sodium hypochlorite lowered the number of colony forming units of *Actinomyces naeslundii* below the detection limit after 10 seconds of irrigation.

Confocal laser scanning microscopy (CLSM) is a reliable method for quantifying bacteria viability and identifying associations between bacterial presence and endodontic infections (29). To quantify bacteria, fluorescent dyes are applied to a biofilm, and they then penetrate the DNA and RNA of bacterial cells. This procedure allows differentiation between living and dead bacterial cells in the root canal system and is useful for studying the antimicrobial efficacy of endodontic treatments before and after instrumentation (30). Given the complexity and bacterial concerns of successful root canal treatment, Neelakantan et al (14) investigated outcomes when using a TEC or TAC design for treatment. They reported debridement of the pulp chambers of mandibular molars was significantly compromised when TAC was performed (14). Because the TAC design retains a truss of dentin between me-

sial and distal access cavities in mandibular molars, the authors were able to evaluate the pulp chambers for reduction in microbial load (14). They concluded that the presence of inaccessible bacteria beneath the truss may affect overall success of endodontic treatment (14). Although many studies have investigated the association between fracture resistance of ETT and different cavity designs (19,20,22,24,31,32), few have investigated microbial reduction of ETT when comparing TEC and TAC (21,24,33). Further, to our knowledge, no studies have investigated TEC and TAC for root canals inoculated with *Actinomyces naeslundii*.

Therefore, the purpose of the current study was to evaluate the impact of access cavity types on the quantity of residual bacteria in the pulp chamber of extracted mandibular first molars after cleaning and shaping during root canal treatment. Specifically, we compared TEC and TAC designs for root canals inoculated with *A. naeslundii* and hypothesized there would be significant differences in the quantity of residual bacteria between designs.

Materials And Methods

Sample Size Analysis

To evaluate the impact of TEC and TAC on residual bacteria after cleaning and shaping during root canal treatment, 30 non-carious mandibular first molars were used. Teeth had been extracted and stored in 10% formalin until use (14). To be included in the study sample, teeth were examined under a high magnification microscope (Global microscope, St. Louis, MO, USA) for cracks or vertical root fractures in mesial or distal roots (5,31). If a tooth had a history of previously initiated root canals or root canal treatment, the presence of internal or external resorptive defects, or severe calcifications, it was excluded from the sample. Once teeth were selected for inclusion, they were evaluated for anatomical similarity. To ver-

Figure 1
TEC and TAC designs used on mandibular first molars and identification of the section used for bacterial quantification.

ify all teeth were as similar as possible, we measured from the occlusal surface to the root apex and measured the mesiodistal and buccolingual dimensions of crowns (32). Teeth with 2 mesial and 1 distal canal were selected based on periapical radiographs. For further standardization of the study sample, additional periapical radiographs were obtained from mesiodistal and buccolingual angles (31,33). The debridement of hard and soft tissues remnants on teeth was manually performed using an ultrasonic scaler. In addition to the 30 teeth for the TEC and TAC groups (15 per each experimental group), we also included 4 controls in the study. Control teeth were prepared using TEC with normal saline as an irrigant, and 2 autoclaved teeth served as negative controls (20,24).

The 30 teeth were randomly divided into the 2 experimental groups. Next, TEC or TAC was performed on the mandibular molars under a high magnification microscope using a size #4 round carbide bur and an Endo Z bur (Dentsply Sirona Endodontics, Ballaigues, Switzerland) with a high-speed

handpiece and copious amounts of water (Figure 1). The TEC was performed according to current guidelines to achieve complete deroofting of the pulp chamber and provide straight-line access to the root canals with no undercuts (12). For TAC, mesial and distal slots of oval cavities were made to gain access over the individual canals on the mesial and distal sides. The roof of the pulp chamber was left intact between the mesial and distal cavities beneath the truss of the tooth (14). Because all teeth in our sample had similar dimensions, we were able to attain some degree of standardization in the size of the access cavities. However, no attempt made to modify the size of the access cavities to gain better access to the canals once cleaning and shaping was initiated. All TEC and TAC procedure were completed by the same endodontics specialist.

The distal canal of the tooth was selected for instrumentation and inoculation of the bacterial sample (Figure 1). The crowns of the teeth were adjusted to have a uniform working length of 18 mm in the distal canals. The apical



Fig. 1A




Fig. 1B

patency of the distal canal was confirmed by using a #10 K-file (Dentsply Sirona Endodontics, Ballaigues, Switzerland) to advance it out of the apical foramen. The canal was gauged up to a #15 K-file, and the working length was determined to be 1 mm short of the apical foramen. All canals were irrigated using 2 mL of 5.25% NaOCl solution performed at 2 mm short of the working length. To neutralize the effect of the sodium hypochlorite, the teeth were immersed in 10% sodium thiosulfate solution for 4 hours followed by a sterile solution for 20 hours. The external surface of the tooth was painted with 2 layers of nail polish to prevent contamination and to seal the apical foramen (23).

Inoculation of Teeth with *Actinomyces naeslundii*

Root canals were inoculated with 100 μ L of an *Actinomyces naeslundii* culture (ATCC 12104 LGC, UK). After inoculation, the bacterial suspension was transferred to the working length of the distal root using a manual #15 K-file. The teeth were then incubated by submersion in Eppendorf tubes containing blood heart infusion broth. The incubation time was 10 days under anaerobic conditions. The culture medium was refreshed every 3 days. The growth of *Actinomyces naeslundii* was confirmed through observation of the turbidity of the solution in the Eppendorf tubes during the incubation period (24).

Root Canal Instrumentation

Teeth were mounted on a metal mounting block, and root canal treatment was performed under a laminar flow hood following strict disinfection protocols. Every treatment was completed by the same individual. All Vortex Blue rotary files (Dentsply Sirona Endodontics, Ballaigues, Switzerland) were attached to a 16:1 gear reduction handpiece (Dentsply, Sirona). Files were used passively to advance into the canal and progress apically. Minimal apical pressure was applied on activation, and 3

easy amplitudes were used in each pass. As suggested by the manufacturer of the Vortex Blue rotary file system, the distal canals in all TEC and TAC teeth were enlarged in a sequence of 15/06, 20/06, 25/06, and 30/06. A standardized volume and duration of irrigant were used for both groups. The canal was irrigated with 1 mL/min of 5.25% NaOCl after every rotary instrument and was recapitulated to working length using a #10 K-file. A hypodermic irrigation needle (30-gauge, side vented, Max-i-Probe, Patterson Dental, USA) was placed passively in the canal and 1 mm short of the apical foramen. The #10 K-file was cleaned with sterile gauze after each recapitulation. A final rinse of 5 mL of 5.25% NaOCl was completed at a rate of 1 mL/min followed by 1 mL/min of rinsing with 17% ethylenediaminetetraacetic acid. An additional rinse was performed using 5 mL/min of sterile saline (34).

The time required to perform the instrumentation and irrigation was recorded and standardized for all (TEC, TAC, and control) groups. Quantification of bacterial load evaluation was performed using CSLM. All teeth were cross-sectioned mesiodistally with a 0.3-mm IsoMet saw at 200 rpm and continuous water. Teeth were stained with BacLight LIVE/DEAD stain (Invitrogen, Carlsbad, CA, USA) for 15 minutes. The CLSM was set at excitation and emission wavelengths of 480 nm and 500 nm. Next, fluorescein diacetate dye was applied, and a 40 \times magnification oil lens was used to evaluate the teeth. The resulting images were analyzed with ImageJ (NIH, LOCI, University of Wisconsin) to quantify live (stained green) and dead (stained red) bacteria (Figure 2). The measured red and green fluorescence intensities were used to calculate the percentage of dead bacteria for both live and dead bacteria (21). Blinding of the operator was not feasible due to the nature of the access cavity designs. However, the examiner responsible for confocal laser scanning microscopy image analysis was blind-

Figure 2
CLSM images of live (green) and dead (red) bacteria in the pulp chambers of teeth using TEC (2A-2B) or TAC (2C-2D) designs for root canal treatment.




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 2D

ed to group allocation.

Statistical Analysis


Bacterial load for all teeth in the TEC and TAC groups was summarized as percentage of bacterial reduction mean and standard deviation (SD). To verify normal distribution of data for percentage of bacterial reduction, a Shapiro-Wilk test was used. For non-normally distributed data, a Wilcoxon-Mann Whitney test was used. For normally distributed data, a 2-sample independent t test was used to evaluate differences in the percentage of bacterial reduction between the TEC and TAC

groups. IBM SPSS Statistics for Windows, Version 27.0 (Armonk, NY; IBM Corp.) statistical software was used for all analyses, and a $P < .05$ was considered significant.

Results

The mean (SD) percentage of bacterial reduction for *Actinomyces naeslundii* was 32.7 (8.0) with a range of 23.3-48.6 for the TEC group and 28.4 (4.8) with a range of 21.6-39.2 for the TAC group. No difference was found between groups ($P = .08$) (Figure 3).

Figure 3
Percentage of bacterial reduction for the TEC and TAC groups.



Discussion

The present in vitro study evaluated the effect of access cavity design on residual bacterial levels in the pulp chamber of mandibular first molars following standardized cleaning and shaping procedures. The results demonstrated no statistically significant difference in the percentage reduction of *Actinomyces naeslundii* between TEC and TAC. Accordingly, the null hypothesis was rejected. These findings are in agreement with previous laboratory investigations reporting comparable bacterial reduction between conventional and conservative access cavity designs when standardized instrumentation and irrigation protocols are employed (21,23,34).

Effective elimination of intracanal microorganisms is essential for periapical healing and long-term success of endodontic treatment (1). Access cavity design plays a pivotal role in facilitating straight-line access, adequate canal instrumentation, and effective irrigant delivery (18). Traditional access cavity preparation, which involves complete deroofing of the pulp chamber, provides unobstructed access to canal orifices and may enhance irrigant penetration (12). However, this approach is associated with substantial loss of coronal and pericervical dentin, which has been shown to compromise fracture resistance and increase susceptibility

to catastrophic failure of endodontically treated teeth (9,13,35).

Fracture remains one of the leading causes of failure in endodontically treated teeth (35). Several studies have demonstrated that conservative access cavity designs, including TAC, improve fracture resistance and result in more favourable fracture patterns compared with TEC (19,32,35). Kishen (35) reported enhanced fracture resistance in teeth prepared with TAC, emphasizing the importance of preserving pericervical dentin. These findings underscore the need to balance structural preservation with effective microbial control to optimize clinical outcomes.

The absence of a significant difference in bacterial reduction between TEC and TAC in the present study suggests that access cavity design alone may have a limited influence on microbial reduction when effective chemomechanical preparation is achieved. Instead, disinfection outcomes appear to be more strongly influenced by instrumentation protocols, irrigant concentration, volume, and contact time. Similar observations were reported by Tüfenkçi and Yılmaz (23), who found no significant difference in *Enterococcus faecalis* reduction between traditional and conservative access cavities. El-Tayeb and Nabeel (21) also demonstrated comparable bacterial reduction between TEC and TAC, while noting inferior disinfection with ultraconservative access cavities.

Studies reporting reduced bacterial reduction with ultraconservative access designs suggest that excessive restriction of coronal access may create coronal interferences, limiting effective instrumentation and irrigant delivery (21,33). In contrast, TAC provides independent access pathways to mesial and distal canals while preserving coronal dentin, potentially minimizing such interferences. Neelakantan et al. (14) reported increased residual pulp tissue beneath the truss in TAC preparations but found no significant differences in canal debridement, supporting the



concept that adequate canal access can still be achieved with TAC designs.

Instrumentation system selection and canal morphology also influence bacterial reduction. Vieira et al. (20) reported increased residual bacteria with conservative access cavities when using XP-Endo Shaper instruments, possibly due to the interaction between access design and instrument kinematics. Conversely, Krishan et al. (19) reported a higher percentage of untouched canal walls in mandibular molars prepared with conservative access cavities, particularly in oval distal canals, highlighting the importance of instrument selection when employing minimally invasive access designs. The choice of *Actinomyces naeslundii* as the test microorganism in the present study was deliberate and clinically relevant. *Actinomyces naeslundii* is a gram-positive, facultative anaerobe frequently associated with persistent primary and secondary endodontic infections and has been implicated in cases of post-treatment apical periodontitis (26–28). Its ability to penetrate dentinal tubules, resist chemomechanical debridement, and survive in harsh intracanal environments makes it a suitable and challenging organism for evaluating disinfection efficacy (26,27). Furthermore, *Actinomyces naeslundii* has been reported to colonize external root surfaces, contributing to persistent periapical pathology even after technically adequate root canal treatment (28). Therefore, its use in this study enhances the clinical relevance of the findings.

Irrigant-related factors also play a critical role in bacterial elimination. Although NaOCl remains the gold standard irrigant, its antimicrobial efficacy is influenced by concentration, volume, and contact time (28,36,37). Lower concentrations may require prolonged exposure to achieve comparable antimicrobial effects. Given the complex anatomy of the root canal system, including isthmuses and accessory canals, adjunctive irrigation

activation techniques such as ultrasonic or laser-assisted irrigation may further enhance disinfection, particularly when conservative access cavity designs are used. Several limitations of this study should be acknowledged. Standardization of canal anatomy was based on periapical radiographs; the use of micro-computed tomography could have provided more precise assessment of canal morphology and reduced anatomical variability. Additionally, only conventional syringe irrigation was evaluated, and adjunctive irrigation activation techniques were not assessed. The in vitro nature of the study also limits direct extrapolation of the findings to clinical conditions. There is a need for future studies focusing on employing novel adjunctives for disinfection (38–40) and also incorporating antimicrobial components into endodontic biomaterials (41).

Within the limitations of this in vitro investigation, bacterial reduction of *Actinomyces naeslundii* was comparable between traditional endodontic and truss access cavity designs following standardized cleaning and shaping procedures. These findings, in conjunction with existing literature (9,21,23), suggest that TAC may be considered a viable minimally invasive alternative to TEC without compromising intracanal disinfection. Further studies incorporating advanced irrigation strategies, guided access techniques, and three-dimensional imaging are warranted to optimize disinfection (42), while preserving tooth structure (43).

Conclusion

Within the limitations of this in vitro study, no statistically significant difference was observed in the reduction of *Actinomyces naeslundii* in the pulp chamber region between TEC and TAC designs following standardized cleaning and shaping procedures. These findings indicate that both access cavity designs provide comparable disinfection of the pulp chamber when ef-



fective chemomechanical preparation is performed. Therefore, TAC may be considered a viable minimally invasive alternative to TEC without compromising bacterial reduction.

Clinical Relevance

Preservation of coronal and pericervical dentin is an important consideration in contemporary endodontic practice. The findings of this study suggest that TAC designs can achieve bacterial reduction comparable to TEC when standard instrumentation and irrigation protocols are used. Accordingly, TAC may be clinically adopted in selected cases to support a minimally invasive endodontic approach while maintaining effective root canal disinfection.

Funding: Nil

Conflicts of Interest: None

References

- Neelakantan P, Romero M, Vera J, Daoud U, Khan AU, Yan A, Cheung GSP. Biofilms in Endodontics-Current Status and Future Directions. *Int J Mol Sci* 2017;18:1748.
- Siqueira JF Jr. Aetiology of root canal treatment failure: why well-treated teeth can fail. *Int Endod J* 2001;34:1-10.
- Tabassum S, Khan FR. Failure of endodontic treatment: The usual suspects. *Eur J Dent* 2016;10:144-147.
- Zadik Y, Sandler V, Bechor R, Salehrabi R. Analysis of factors related to extraction of endodontically treated teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:31-5.
- Teja KV, Janani K, Alqahtani AA, Robaian A, Alhalabi F, Merdad KA, Alam MK, Shrivastava D, Jose J, Srivastava KC. Herbal Agents versus Ethylene Diamine Tetra Acetic Acid on Removal of the Smear Layer-A Systematic Review of In Vitro Studies. *Int J Environ Res Public Health* 2022;19:6870.
- Teja KV, Vasundhara KA, Georgiou AC, Spagnuolo G, Esposito L, Armogida NG. Efficacy of Gel- and Solution-Based Formulations of Conventional Endodontic Disinfectants During Root Canal Shaping: A Systematic Review of Laboratory and Clinical Studies. *Eur Endod J*. 2025 ;10:447-455.
- Alghamdi F, Shakir M. The Influence of *Enterococcus faecalis* as a Dental Root Canal Pathogen on Endodontic Treatment: A Systematic Review. *Cureus* 2020;12:7257.
- Peters Hayati AT, Prisinda D, Nugroho ALL. Survival and Success Rate of Restoration Post Endodontic Treatment. *Clin Cosmet Investig Dent* 2025;17:525-537.
- Versiani MA, Kang S, Kim HC, Cho M, Silva EJNL, Belladonna FG, et al. Influence of Soffit Preservation on Root Stress Distribution in Mandibular Molars: A Finite Element Study. *J Endod* 2025;51:1625-1636.
- Gluskin AH, Peters CI, Peters OA. Minimally invasive endodontics: challenging prevailing paradigms. *Br Dent J* 2014;216:347-53.
- Silva EJNL, Pinto KP, Ferreira CM, Belladonna FG, De-Deus G, Dummer PMH, Versiani MA. Current status on minimal access cavity preparations: a critical analysis and a proposal for a universal nomenclature. *Int Endod J* 2020;53:1618-1635.
- Patel S, Rhodes J. A practical guide to endodontic access cavity preparation in molar teeth. *Br Dent J*. 2007 Aug 11;203(3):133-40.
- Clark D, Khademi JA. Case studies in modern molar endodontic access and directed dentin conservation. *Dent Clin North Am* 2010;54:275-89.
- Neelakantan P, Khan K, Hei Ng GP, Yip CY, Zhang C, Pan Cheung GS. Does the Orifice-directed Dentin Conservation Access Design Debride Pulp Chamber and Mesial Root Canal Systems of Mandibular Molars Similar to a Traditional Access Design? *J Endod* 2018;44:274-279.
- Selvakumar RJ, Surendran S, Sundar S, Arul B, Natanasabapathy V. Impact of Contracted Endodontic Access Cavities on the Fracture Resistance of Endodontically Treated Teeth After Mechanical Aging by Simulated Chewing Forces. *J Endod* 2023;49:1176-1182.
- Torbjörner A, Fransson B. A literature review on the prosthetic treatment of structurally compromised teeth. *Int J Prosthodont* 2004;17:369-76.
- Peutzfeldt A, Sahafi A, Asmussen E. A survey of failed post-retained restorations. *Clin Oral Investig* 2008;12:37-44.
- Silva AA, Belladonna FG, Rover G, Lopes RT, Moreira E JL, De-Deus G, Silva EJNL. Does ultra-conservative access affect the efficacy of root canal treatment and the fracture resistance of two-rooted maxillary premolars? *Int Endod J* 2020;53:265-275.
- Krishan R, Paqué F, Ossareh A, Kishen A, Dao T, Friedman S. Impacts of conservative endodontic cavity on root canal instrumentation efficacy and resistance to fracture assessed in incisors, premolars, and molars. *J Endod* 2014;40:1160-6.
- Vieira GCS, Pérez AR, Alves FRF, Provenzano JC, Mdala I, Siqueira JF Jr, et al. Impact of Contracted Endodontic Cavities on Root Canal Disinfection and Shaping. *J Endod* 2020;46:655-661.
- El-Tayeb MMN, Nabeel M. Impact of contracted endodontic access cavity designs on root canal disinfection using diode laser: an "in-vitro" study. *Int Arab J Dent* 2023;14:130-141.
- Ivanoff CS, Marchesan MA, Andonov B, Hottel TL, Dandarov Y, Mandova S, Iftikhar H. Fracture resistance of mandibular premolars with contracted or traditional endodontic access cavities and class II temporary composite restorations. *Endod Pract Today* 2017;11:7-14.
- Tüfenkçi P, Yılmaz K. The Effects of Different Endodontic Access Cavity Design and Using XP-endo Finisher on the Reduction of *Enterococcus fae-*

- calis in the Root Canal System. *J Endod* 2020;46:419-424.
24. Barbosa AFA, Silva EJNL, Coelho BP, Ferreira CMA, Lima CO, Sassone LM. The influence of endodontic access cavity design on the efficacy of canal instrumentation, microbial reduction, root canal filling and fracture resistance in mandibular molars. *Int Endod J* 2020;53:1666-1679.
 25. Radcliffe CE, Potouridou L, Qureshi R, Hababeh N, Qualtrough A, Worthington H, et al. Antimicrobial activity of varying concentrations of sodium hypochlorite on the endodontic microorganisms *Actinomyces israelii*, *A. naeslundii*, *Candida albicans* and *Enterococcus faecalis*. *Int Endod J* 2004;37:438-46.
 26. Sunde PT, Olsen I, Debelian GJ, Tronstad L. Microbiota of periapical lesions refractory to endodontic therapy. *J Endod* 2002;28:304-10.
 27. Dioguardi M, Quarta C, Alovisei M, Crincoli V, Aiuto R, Crippa R, et al. Microbial Association with Genus *Actinomyces* in Primary and Secondary Endodontic Lesions, Review. *Antibiotics* 2020;9:433.
 28. Esteves LS, Henriques ACG, Silva CAVME, Cangussu MCT, Ramos EAG, Estrela C, Santos JND. Actinomycosis is not Frequent in the Periapex But is a Persistent Lesion. *Braz Dent J* 2017;28:688-693.
 29. Mattern R, Ernst S, Böcher S, Braun A, Wenzler JS, Conrads G. CLSM-Guided Imaging for Quantifying Endodontic Disinfection. *Antibiotics* 2024;13:54.
 30. Carvalho FG, Puppim-Rontani RM, Fúcio SB, Negrini Tde C, Carlo HL, Garcia-Godoy F. Analysis by confocal laser scanning microscopy of the MDPB bactericidal effect on *S. mutans* biofilm CLSM analysis of MDPB bactericidal effect on biofilm. *J Appl Oral Sci* 2012;20:568-75.
 31. Daniel A, Saleh AR, Al-Jadaa A, Kheder W. Impact of Access Cavity Design on Fracture Resistance of Endodontically Treated Maxillary First Premolar: In Vitro. *Braz Dent J* 2024;35:245676.
 32. Özyürek T, Ülker Ö, Demiryürek EÖ, Yılmaz F. The Effects of Endodontic Access Cavity Preparation Design on the Fracture Strength of Endodontically Treated Teeth: Traditional Versus Conservative Preparation. *J Endod* 2018;44:800-805.
 33. Andac G, Kalender A, Baddal B, Basmaci F. Impact of different access cavity designs and Ni-Ti files on the elimination of enterococcus faecalis from the root canal system: An In Vitro Study. *Appl Sci* 2022;12:2049.
 34. Kandaswamy D, Venkateshbabu N. Root canal irrigants. *J Conserv Dent* 2010;13:256-64.
 35. Kishen A. Mechanisms and risk factors for fracture predilection in endodontically treated teeth. *Endod Topics* 2006;13:57-83.
 36. Byström A, Immich F, Rödiger T, Kanzow P, Piva E, Rossi-Fedele G. Effectiveness of Root Canal Irrigation and Dressing for the Treatment of Asymptomatic Apical Periodontitis: A Systematic Review Assessing Complementary Timeframes. *Aust Endod J* 2025;51:209-217.
 37. Prasad N, Dasson Bajaj P, Shenoy R, Dutta A, Thomas MS. Sodium Hypochlorite Concentration and Postendodontic Pain - Unveiling the Optimal Balance: A Systematic Review and Meta-Analysis. *J Endod* 2024;50:1233-1244.
 38. Rajamanickam K, Teja KV, Ramesh S, Choudhari S, Cernera M, Armogida NG, Mustafa M, Spagnuolo G. Evaluation of root canal cleanliness on using a novel irrigation device with an ultrasonic activation technique: an ex vivo study. *Applied Sci* 2023 Jan 6;13:796.
 39. Penukonda R, Teja KV, Kacharaju KR, Xuan SY, Sheun LY, Cernera M, Iaculli F. Comparative evaluation of smear layer removal with Ultra-X device and XP-Endo Finisher file system: An ex-vivo study: Smear removal on using various activation devices. *Giornale Italiano di Endod* 2023;37(2).
 40. Rajamanickam K, Teja KV, Ramesh S, AbuMelha AS, Alkahtany MF, Almadi KH, Bahammam SA, Janani K, Choudhari S, Jose J, Srivastava KC, Shrivastava D, Patil S. Comparative Study Assessing the Canal Cleanliness Using Automated Device and Conventional Syringe Needle for Root Canal Irrigation-An Ex-Vivo Study. *Materials* 2022;15:6184.
 41. Jose J, Teja KV, Janani K, Alam MK, Khattak O, Salloum MG, Magar SS, Magar SP, Rajeshkumar S, Palanivelu A, Srivastava KC, Shrivastava D. Preparation of a Novel Nanocomposite and Its Antibacterial Effectiveness against *Enterococcus faecalis*-An In Vitro Evaluation. *Polymers* 2022;14:1499.
 42. Iandolo A, Armogida NG, Mancino D, Spagnuolo G, Cernera M, Abdellatif D. Evaluation of Root Canal Cleaning and Irrigant Penetration Using Different Irrigation Protocols: A Combined SEM and Single-Tooth Micro-CT Study. *Clin Exp Dent Res* 2025;11:e70175.
 43. Spinelli V, Gallicchio V, Traino F, Marino C, De Santis R, Rengo S, Armogida NG. Influence of the direct composite restoration on the fracture strength and failure modes of endodontically treated premolars restored with posts. *Giornale Italiano di Endod* 2025;39(1).